

February 1, 2008

Diffeomorphisms of Curved Manifolds

J.S. Apps

Department of Theoretical Physics
The University, Manchester M13 9PL

ABSTRACT

We obtain an expression for the curvature of the Lie group $\text{SDiff}\mathcal{M}$ and use it to derive Lukatskii's formula for the case where $\mathcal{M}$ is locally Euclidean. We discuss qualitatively some previous findings for $\text{SDiff}S^2$ in conjunction with our result.

1 Preliminaries

We consider the Lie group $\text{SDiff}\mathcal{M}$ of volume-preserving diffeomorphisms of a d -dimensional closed Riemannian manifold $\mathcal{M}$. Details are in Arnold [1]. The Lie algebra is the set of solenoidal vector fields on $\mathcal{M}$, endowed with the metric

$$\langle U, V \rangle = \int_{\mathcal{M}} d^d x \sqrt{g} \, U \cdot V \quad (1)$$

If the metric and fields are invariant under the right-action of the diffeomorphism group, then we require the covariant derivative on $\text{SDiff}\mathcal{M}$ to satisfy

$$\nabla_U \langle V, W \rangle = 0 \quad (2)$$

so that, if the metric on $\text{SDiff}\mathcal{M}$ is Riemannian,

$$\langle \nabla_U V, W \rangle + \langle V, \nabla_U W \rangle = 0 \quad (3)$$

We define D_U to be the covariant derivative on $\mathcal{M}$ in the direction of the field U . Since

$$(D_U V) \cdot W + V \cdot (D_U W) = D_U(V \cdot W) = U^\mu \partial_\mu (V \cdot W) \quad (4)$$

we have

$$\langle D_U V, W \rangle + \langle V, D_U W \rangle = \int_{\mathcal{M}} d^d x \sqrt{g} \, U^\mu \partial_\mu (V \cdot W) = 0 \quad (5)$$

where the latter equality follows from integration by parts, together with the solenoidal condition $\partial_\mu(\sqrt{g} \, U^\mu) = 0$. Note that (5) extends to the case where V and W have a non-zero divergence.

We make the orthogonal decomposition

$$D_U V = P_U V + Q_U V \quad (6)$$

where $P_U V$ is the solenoidal projection, and, from (5), satisfies

$$\langle P_U V, W \rangle + \langle V, P_U W \rangle = 0 \quad (7)$$

for V and W solenoidal. $P_U V$ is therefore a vector on $\text{SDiff}\mathcal{M}$ which satisfies the criterion (3) for a covariant derivative of right-invariant fields. Additionally, since the Lie bracket $[U, V] = D_U V - D_V U$ of two solenoidal fields U and V is itself solenoidal, we have

$$P_U V - P_V U = [U, V] \quad (8)$$

so that P is the torsionless connection on $\text{SDiff}\mathcal{M}$, i.e. $P_U V = \nabla_U V$. The curvature of $\text{SDiff}\mathcal{M}$ with respect to fields U, V, W and X is then defined:

$$T_{UVWX} = \langle (P_{[U,V]} W - [P_U, P_V] W), X \rangle \quad (9)$$

We write this as

$$\langle [D_{[U,V]} W - D_U(D_V W - Q_V W) + D_V(D_U W - Q_U W)], X \rangle$$

since X is orthogonal to the Q projection of D . Using (5), this becomes

$$\langle (D_{[U,V]} W - [D_U, D_V] W), X \rangle - \langle Q_V W, Q_U X \rangle + \langle Q_U W, Q_V X \rangle$$

The first term is simply

$$\int_{\mathcal{M}} d^d x \sqrt{g} U^\mu V^\nu W^\rho X^\sigma R_{\mu\nu\rho\sigma}$$

where $R_{\mu\nu\rho\sigma}$ are the components of the Riemann tensor on some coordinate basis on $\mathcal{M}$.

2 Locally Euclidean Manifolds

If $\mathcal{M}$ is locally Euclidean, then $R_{\mu\nu\rho\sigma} = 0$. As $Q_U V = Q_V U$, we obtain Lukatskii's formula [2] for the (unnormalized) sectional curvature T_{UVUV} :

$$T_{UVUV} = \frac{1}{4} [\langle \hat{\chi}(U, U), \hat{\chi}(V, V) \rangle - \langle \hat{\chi}(U, V), \hat{\chi}(U, V) \rangle] \quad (10)$$

where $\chi(U, V) = D_U V + D_V U$, and the hat denotes the projection orthogonal to $\text{SDiff}\mathcal{M}$. Note that we suspect a typing error in equation (3) of [2], which we have corrected: the “+” should be replaced by a “−”.

3 The General Case

In all cases we can express $Q_U V$ as the gradient of a single-valued scalar field (see Arnold, p 341):

$$Q_U V = \text{grad } \xi \Rightarrow \text{div } (D_U V) = \Delta \xi \quad (11)$$

Writing $D_U = U \cdot \text{grad}$ and integrating by parts, our result is

$$\begin{aligned} T_{UVUV} &= \int_{\mathcal{M}} d^d x \sqrt{g} \{ U^\mu V^\nu U^\rho V^\sigma R_{\mu\nu\rho\sigma} \\ &+ \text{div } [(U \cdot \text{grad})V] \Delta^{-1} \text{div } [(U \cdot \text{grad})V] \\ &- \text{div } [(U \cdot \text{grad})U] \Delta^{-1} \text{div } [(V \cdot \text{grad})V] \} \end{aligned} \quad (12)$$

The sectional curvature of $\mathcal{M}$ in the plane defined by the vectors U and V is given by

$$K_{\mathcal{M}}(U, V) = \frac{U^\mu V^\nu U^\rho V^\sigma R_{\mu\nu\rho\sigma}}{U^\mu V^\nu U^\rho V^\sigma (g_{\mu\rho} g_{\nu\sigma} - g_{\mu\sigma} g_{\nu\rho})} \quad (13)$$

so that the first term in (12) is

$$\int_{\mathcal{M}} d^d x \sqrt{g} [(U \cdot U)(V \cdot V) - (U \cdot V)^2] K_{\mathcal{M}}(U, V)$$

Since $(U \cdot U)(V \cdot V) - (U \cdot V)^2 \geq 0$, the sign of this term is definite if the sign of $K_{\mathcal{M}}(U, V)$ is definite.

In particular, for the case $d = 2$, $K_{\mathcal{M}}(U, V)$ is independent of U and V , and is equal to $R/2$, where R is the Ricci scalar. Also, a solenoidal field on $\mathcal{M}$ is the curl of an (in general multi-valued) scalar potential :

$$U^\mu = \frac{\epsilon^{\mu\nu}}{\sqrt{g}} \partial_\nu \chi \quad (14)$$

If the potentials for U and V are χ and ψ respectively then the first term in (12) becomes

$$\int_{\mathcal{M}} d^2 x \sqrt{g} \frac{1}{2} R \{\chi, \psi\}^2$$

where $\{\chi, \psi\}$ is the generally covariant Poisson bracket $(\epsilon^{\mu\nu} \partial_\nu \chi \partial_\mu \psi) / \sqrt{g}$.

4 The 2-sphere

As an example, we calculate the sectional curvature of $\text{SDiff}S^2$, alternative procedures for which have already been established by Arakelyan and Savvidy [3] and Lukatskii [4]. The sphere is of unit radius, so that $R = 2$. We choose the diffeomorphism generators $U = \text{curl}(\cos \theta)$ and $V = \text{curl}[\sin \theta(e^{i\phi} + e^{-i\phi})]$. For this simple case, all relevant quantities turn out to be proportional to spherical harmonics, so calculation of the inverse Laplacian is easy.

$$\begin{aligned}
\text{div}[(U \cdot \text{grad})U] &= 1 - 3\cos^2 \theta \\
\text{div}[(V \cdot \text{grad})V] &= 2(3\cos^2 \theta - 1) \\
&\Rightarrow \Delta^{-1} \text{div}[(V \cdot \text{grad})V] = -\frac{1}{3}(3\cos^2 \theta - 1) \\
\text{div}[(U \cdot \text{grad})V] &= -3\sin \theta \cos \theta(e^{i\phi} + e^{-i\phi}) \\
&\Rightarrow \Delta^{-1} \text{div}[(U \cdot \text{grad})V] = \frac{1}{2}\sin \theta \cos \theta(e^{i\phi} + e^{-i\phi})
\end{aligned}$$

We obtain

$$\int_{S^2} d^2x \sqrt{g} \frac{1}{2} R \{\chi, \psi\}^2 = \frac{16\pi}{3} \quad (15)$$

$$\int_{S^2} d^2x \sqrt{g} \text{div}[(U \cdot \text{grad})V] \Delta^{-1} \text{div}[(U \cdot \text{grad})V] = -\frac{8\pi}{5} \quad (16)$$

$$\int_{S^2} d^2x \sqrt{g} \text{div}[(U \cdot \text{grad})U] \Delta^{-1} \text{div}[(V \cdot \text{grad})V] = \frac{16\pi}{15} \quad (17)$$

As $\langle U, U \rangle = 8\pi/3$, $\langle V, V \rangle = 32\pi/3$, $\langle U, V \rangle = 0$, the normalized sectional curvature is

$$K_S(U, V) = \frac{\frac{16\pi}{3} - \frac{8\pi}{5} - \frac{16\pi}{15}}{\frac{8\pi}{3} \times \frac{32\pi}{3}} = \frac{3}{32\pi} \quad (18)$$

This agrees with the result found by Lukatskii [4] and by Dowker and Mo-zheng [5] for the same vectors. In this instance, our method is much more tedious than the alternative, but this may not be true for a more general case, in which summations over spherical harmonics are involved.

5 Discussion

The theory of volume-preserving diffeomorphism groups may be applied to fluid dynamics, where the diffeomorphism generators give the velocity field of an incompressible fluid on $\mathcal{M}$ (see [1]). Given two flows with initial velocity fields U and V , a negative value of $K_S(U, V)$ causes the difference in the fluid configurations to diverge with time. In this way, the “stability” of a flow can be predicted.

We may construct a very heuristic argument concerning the origin of the individual terms in (12), with reference to the fluid dynamical interpretation. Since the second and third terms arise from taking the solenoidal projection of the covariant derivative, they can be seen as contributions due to forces between fluid elements which maintain the volume-preserving nature of the flow. The first term, on the other hand, only arises when $\mathcal{M}$ is curved, and may be interpreted as an effect on the flow stability due to the deviation of geodesics on $\mathcal{M}$. We therefore write

$$K_S(U, V) = K_C(U, V) + K_F(U, V) \quad (19)$$

where $K_C(U, V)$ is the “curvature” term.

In the case of the sphere, there was some surprise (see [5]) that the sectional curvature $K_S(U, V)$ for $U = \text{curl}(\cos \theta)$ turned out to be positive, since for similar flows on a torus, the value is negative [1]. In the light of the preceding argument, it seems more appropriate to compare only $K_F(U, V)$ with Arnold’s results, since the torus is flat.

We restrict ourselves to the case

$$\begin{aligned} U &= \text{curl}(\cos \theta) \\ V &= \text{curl}[Y_l^m(\theta, \phi) + (-1)^m Y_l^{-m}(\theta, \phi)] \end{aligned}$$

so that U and V are real. It is easy to show that

$$\{\cos \theta, Y_l^m(\theta, \phi)\} = imY_l^m(\theta, \phi)$$

and then use this to derive

$$K_C(U, V) = \frac{3m^2}{8\pi l(l+1)} \quad (20)$$

The formula in [4] and [5]:

$$K_S(U, V) = \frac{3m^2}{8\pi[l(l+1)]^2} \quad (21)$$

then gives us

$$K_F(U, V) = \frac{3m^2}{8\pi l(l+1)} \left[\frac{1}{l(l+1)} - 1 \right] \quad (22)$$

so that K_F is always negative. We suggest that the positive curvature of the sphere “stabilizes” the flow.

In general, we hope that equation (12) will facilitate the calculation of the curvature of $\text{SDiff}\mathcal{M}$ for other manifolds. This is certainly true for the torus: the use of Lukatskii’s formula saves a great deal of work in deriving Arnold’s expression for the curvature.

I would like to thank Stuart Dowker for helpful discussions.

References

- [1] V.I. Arnold, Ann. Inst. Fourier **16**, 319 (1966).
- [2] A.M. Lukatskii, Russian Mathematical Surveys **45**, 160 (1990).
- [3] T.A. Arakelyan and G.K. Savvidy, Phys. Lett. B **223**, 41 (1989)
- [4] A.M. Lukatskii, Funct. Anal. Appl. **13**, 174 (1980)
- [5] J.S. Dowker and Wei Mo-zheng, Class. Quantum Grav. **7**, 2361 (1990).